

REGULARIZATION OF CURRENT ALGEBRA

JOUKO MICKELSSON

Department of Mathematics, University of Jyväskylä, SF-40100, Finland

ABSTRACT In this talk I want to explain the operator subtractions needed to regularize gauge currents in a second quantized theory. The case of space-time dimensions $3 + 1$ is considered in detail. In presence of chiral fermions the regularization effects a modification of the local commutation relations of the currents by local Schwinger terms. In $1 + 1$ dimensions one gets the usual central extension (Schwinger term does not depend on background gauge field) whereas in $3 + 1$ dimensions one gets an anomaly linear in the background potential.

1. INTRODUCTION

Chiral fermions in a nonabelian external gauge field are quantized as follows. Let G be a compact gauge group, $\mathfrak{g}$ its Lie algebra, M the physical space, and $\mathcal{A}$ the space of smooth $\mathfrak{g}$ valued vector potentials in M . For each $A \in \mathcal{A}$ one constructs a fermionic Fock space $\mathcal{F}_A$ containing a Dirac vacuum ψ_A . The Hilbert space $\mathcal{F}_A$ carries an irreducible representation of the canonical anticommutation relations

Presented at the conference "Constraint Theory and Quantization" in Montepulciano, June 1993

(CAR)

$$a^*(u)a(v) + a(v)a^*(u) = (u, v) \text{ all other anticommutators} = 0.$$

The representation is characterized by the property

$$(1.1) \quad a^*(u)\psi_A = 0 = a(v)\psi_A \text{ for } u \in H_-(A) \text{ and } v \in H_+(A)$$

where $H_+(A)$ is the subspace of the one-particle fermionic Hilbert space H spanned by the eigenvectors of the Dirac-Weyl Hamiltonian

$$(1.2) \quad D_A = i\gamma_k(\nabla_k + A_k)$$

belonging to nonnegative eigenvalues and $H_-(A)$ is the orthogonal complement of $H_+(A)$. Here ∇_k 's are covariant derivatives in directions given by a (local) orthonormal basis, with respect to a fixed Riemannian metric on M . In the following we shall concentrate to the physically most interesting case $\dim M = 3$ and the γ -matrices can be chosen as the Pauli matrices $\sigma_1, \sigma_2, \sigma_3$ with $\sigma_1\sigma_2 = i\sigma_3$ (and similarly for cyclic permutations of the indices) and $\sigma_k^2 = 1$.

The group $\mathcal{G} = \text{Map}(M, G)$ of smooth gauge transformations acts on $\mathcal{A}$ as $g \cdot A = gAg^{-1} + dg g^{-1}$. The Fock spaces $\mathcal{F}_A$ form a vector bundle over $\mathcal{A}$. A natural question is then: How does $\mathcal{G}$ act in the total space $\mathcal{F}$ of the vector bundle? Since the base $\mathcal{A}$ is flat there obviously is a lift of the action on the base to the total space. However, we have the additional physical requirement that

$$(1.3) \quad \hat{g}\hat{D}_A\hat{g}^{-1} = \hat{D}_{g \cdot A}$$

where $\hat{D}_A$ is the second quantized Hamiltonian and $\hat{g}$ is the lift of g to $\mathcal{F}$. This condition has as a consequence that $\hat{g}\psi_A$ should be equal, up to a phase, to the vacuum ψ_0 .

A complication in all space-time dimensions higher than $1 + 1$ is that the representations of CAR in the different fibers of $\mathcal{F}$ are inequivalent, [A]. The effect of this is that a proper mathematical definition of the infinitesimal generators of $\mathcal{G}$ (current algebra) involves further renormalizations in addition to the normal ordering prescription. In one space dimensions the situation is simple. The current algebra is contained in a Lie algebra $\mathbf{gl}_1$ which by definition consists of all bounded operators X in H satisfying $[\epsilon, X] \in L_2$, where ϵ is the sign operator $\frac{D_0}{|D_0|}$ associated to the free Dirac operator and L_2 is the space of Hilbert-Schmidt operators. In general, we denote by L_p the Schatten ideal of operators T with $|T|^p$ a trace-class operator. Let $a_n^* = a^*(u_n)$, where $D_0 u_n = \lambda_n u_n$ and the eigenvalues are indexed such that $\lambda_n \geq 0$ for $n \geq 0$ and $\lambda_n < 0$ for $n < 0$. Denoting the matrix elements of a one-particle operator X by (X_{nm}) , the second quantized operator $\hat{X}$ is

$$(1.4) \quad \hat{X} = \sum X_{nm} : a_n^* a_m :$$

where the normal ordering is defined by

$$: a_n^* a_m := \begin{cases} a_m a_n^* & \text{if } n = m < 0 \\ a_n^* a_m & \text{otherwise} . \end{cases}$$

The commutation relations are

$$(1.5) \quad [\hat{X}, \hat{Y}] = \widehat{[X, Y]} + c(X, Y)$$

where c is the Lundberg's cocycle, [L],

$$(1.6) \quad c(X, Y) = \frac{1}{4} \text{tr} \epsilon [\epsilon, X] [\epsilon, Y].$$

When X, Y are infinitesimal gauge transformations on a circle the right-hand-side is equal to the central term of an affine Kac-Moody algebra, [PS],

$$(1.7) \quad c(X, Y) = \frac{i}{4\pi} \int \text{tr} X' Y.$$

In this talk I want to explain the regularizations needed in $3+1$ space-time dimensions and the generalization of (1.4) through (1.7).

2. ACTION OF THE GROUP OF GAUGE TRANSFORMATIONS IN THE FOCK BUNDLE

Let $\epsilon(A) = \frac{D_A}{|D_A|}$; if D_A has zero modes define $\epsilon(A)$ to be $+1$ in the zero mode subspace. For $A \in \mathcal{A}$ denote by P_A the set of unitary operators $h : H \rightarrow H$ such that

$$(2.1) \quad [\epsilon, h^{-1}\epsilon(A)h] \in L_2.$$

If $h \in P_A$ then also $hs \in P_A$ for any $s \in U_1$, where U_1 is the group of unitary operators s with the property $[\epsilon, s] \in L_2$. The spaces P_A form a principal bundle over $\mathcal{A}$ with the structure group U_1 .

Since $\mathcal{A}$ is flat the bundle P is trivial and we may choose a section $A \mapsto h_A \in P_A$. Define

$$(2.2) \quad \omega(g; A) = h_{g \cdot A}^{-1} T(g) h_A$$

where $T(g)$ is the one-particle representation of $g \in \mathcal{G}$. By construction, ω satisfies the 1-cocycle condition

$$(2.3) \quad \omega(gg'; A) = \omega(g; g' \cdot A) \omega(g'; A).$$

Using $T(g)D_A T(g)^{-1} = D_{g \cdot A}$ we get $T(g)\epsilon(A)T(g)^{-1} = \epsilon(g \cdot A)$ which implies

$$h_{g \cdot A}[\epsilon, \omega(g; A)]h_A^{-1} = (h_{g \cdot A}\epsilon h_{g \cdot A}^{-1})T(g) - T(g)(h_A\epsilon h_A^{-1})$$

$$= \epsilon(g \cdot A)T(g) - T(g)\epsilon(A) \bmod L_2 = 0$$

Since L_2 is an operator ideal this equation implies

$$(2.4) \quad [\epsilon, \omega(g; A)] \in L_2.$$

Thus the 1-cocycle ω takes values in the group U_1 .

The group valued cocycle ω gives rise to a Lie algebra cocycle θ by

$$(2.5) \quad \begin{aligned} \theta(X; A) &= \frac{d}{dt} \omega(e^{tX}; A)|_{t=0} \\ &= h_A^{-1} dT(X) h_A - h_A^{-1} \mathcal{L}_X h_A. \end{aligned}$$

It satisfies the Lie algebra cocycle condition

$$(2.6) \quad \theta([X, Y]; A) - [\theta(X; A), \theta(Y; A)] + \mathcal{L}_X \theta(Y; A) - \mathcal{L}_Y \theta(X; A) = 0,$$

where $\mathcal{L}_X$ is the Lie derivative in the direction of the infinitesimal gauge transformation X , $\mathcal{L}_X f(A) = \frac{d}{dt} f(e^{tX} \cdot A)|_{t=0}$. We denote by dT the Lie algebra representation in H corresponding to the representation T of finite gauge transformations. For each $A \in \mathcal{A}$ and $X \in \text{Map}(M, \mathfrak{g})$ the operator $\theta(X; A) \in \mathfrak{gl}_1$.

The section h_A of P can be used to trivialize the bundle of Fock spaces over $\mathcal{A}$. Each fiber $\mathcal{F}_A$ is identified as the free Fock space $\mathcal{F}_0$. The Hamiltonian D_A is quantized as

$$(2.7) \quad \hat{D}_A = q(h_A^{-1} D_A h_A),$$

that is, we first conjugate the one-particle operator D_A by h_A and then canonically quantize $h_A^{-1} D_A h_A$. The conjugated operator has a Dirac vacuum ψ_A contained in F_0 (but differing from the free vacuum ψ_0). The CAR algebra in the background A is represented in $\mathcal{F}_0$ through the automorphism $a^*(u) \mapsto a_A^*(u) = a^*(h_A^{-1}u)$, $a(u) \mapsto a_A(u) = a(h_A^{-1}u)$ and using the free CAR representation for the operators on the right. The Hamiltonian $\hat{D}_A$ is then

$$(2.8) \quad \hat{D}_A = \sum : a_A^*(y_+) a_A(y_-) :$$

where the u_n 's for nonnegative (negative) indices are the eigenvectors of D_A belonging to nonnegative (negative) eigenvalues. The normal ordering is defined with respect to the free vacuum.

Sections of the Fock bundle are now ordinary $\mathcal{F}_0$ valued functions. The effect of an infinitesimal gauge transformation consists of two parts: The Lie derivative $\mathcal{L}_X$ acting on the argument A of the function and an operator acting in $\mathcal{F}_0$,

$$(2.9) \quad \hat{X} = \mathcal{L}_X + \sum \theta(X; A)_{nm} : a_n^* a_m :,$$

where the $\theta(X; A)_{nm}$'s are matrix elements of $\theta(X; A)$ in the eigenvector basis (v_n) of D_0 . The commutation relations of the second quantized operators are modified by the Lundberg's cocycle, [M1],

$$(2.10) \quad [\hat{X}, \hat{Y}] = \widehat{[X, Y]} + c(\theta(X; A), \theta(Y; A)).$$

In the next section we want to compute the right-hand side of (2.10) more explicitly.

We shall denote by $c_n(X, Y; A)$ ($n = \dim M$) the second term on the right. It is a Lie algebra 2-cocycle in the following sense:

$$c_n([X, Y], Z; A) + \mathcal{L}_X c_n(Y, Z; A) + \text{cyclic perm.} = 0.$$

3. A COMPUTATION OF THE COCYCLE

First we shall construct the section h_A explicitly as a function of the vector potential when $\dim M = 3$. We shall define h_A through its symbol, as a pseudo-differential operator (PSDO) in the spin bundle over M . I claim that an operator with the following asymptotic expansion satisfies the requirement (2.1):

$$(3.1) \quad h_A = 1 + \frac{i}{4} \frac{[\xi, A]}{|\xi|^2} + \text{terms of lower order in } |\xi|.$$

Here $\xi = \sum \xi_k \sigma_k$ is the three-momentum; its components represent partial derivatives $-i\partial_k$ in M , with respect to some local coordinates. In order to make the discussion as simple as possible we assume that M is the one-point compactification of $\mathbb{R}^3$ and we use standard coordinates in $\mathbb{R}^3$. We also use the notation $A = \sum A_k \sigma_k$.

An example of an unitary operator with the asymptotic expansion (3.1) is the operator

$$(3.2) \quad h_A = \exp\left(\frac{i}{4}(D_0^2 + \lambda)^{-1/2}[D_0, A](D_0^2 + \lambda)^{-1/2}\right)$$

where we have added a small positive constant λ to the denominator in order to cancel the infrared singularity at $\xi = 0$; this has an effect in the asymptotic expansion only on terms of order -2 and lower in the momentum ξ . It is clear that the lower order terms do not have any effect on the condition (2.1) since any operator of order ≤ -2 is automatically Hilbert-Schmidt when the dimension of M is 3. Thus we have

$$(3.3) \quad \theta(X; A) = h_A^{-1} dT(X) h_A - h_A^{-1} \mathcal{L}_X h_A = X + \frac{i}{4} \frac{[\xi, A]}{|\xi|^2} + O(-2)$$

where $O(-p)$ denotes terms of order $\leq -p$. The symbol of the PSDO ϵ is $\frac{\xi}{|\xi|}$ and it is a simple computation to check that indeed $[\epsilon, \theta(X; A)] \in L_2$ using the product rule of symbols,

$$(3.4) \quad (p * q)(\xi, x) = \sum_n \frac{(-i)^{|n|}}{n!} \partial_\xi^n p \partial_x^n q$$

where the sum is over multi-indices $n = (n_1, n_2, n_3) \in \mathbb{N}^3$, $|n| = n_1 + n_2 + n_3$, $n! = n_1! n_2! n_3!$ and $\partial_x^n = (\frac{\partial}{\partial x_1})^{n_1} (\frac{\partial}{\partial x_2})^{n_2} (\frac{\partial}{\partial x_3})^{n_3}$.

The term of order -2 in θ is important in computing the actual value of θ . It is equal to

$$(3.5) \quad \begin{aligned} \theta_{-2} = & -\frac{1}{4} \frac{[\sigma_k, A]}{|\xi|^2} \partial_k X + \frac{1}{2} \frac{[\xi, A]}{|\xi|^4} \xi_k \partial_k X \\ & + \frac{1}{16} \frac{[\xi, A]}{|\xi|^4} [\xi, dX]. \end{aligned}$$

Note that all terms are linear in the vector potential A . The computation of $c_3(X, Y; A) = c(\theta(X; A), \theta(Y; A))$ is greatly simplified when we keep in mind that it is only the cohomology class of the cocycle c_3 we are interested in. Another simplification is the following: Formally,

$$(3.6) \quad \frac{1}{4} \text{tr} \epsilon[\epsilon, P][\epsilon, Q] = -\frac{1}{2} \text{tr}[\epsilon, P]Q$$

when P, Q are in $\mathfrak{gl}_1$. However, the operator on the right is not quite trace-class; only its diagonal blocks are trace-class. For this reason the trace is only conditionally convergent. It is convergent when evaluated with respect to a basis compatible with the polarization $H = H_+ \oplus H_-$, for example, one can choose a basis of eigenvectors of D_0 . The trace of an operator P with symbol $p(\xi, x)$ on a n -dimensional manifold is

$$(3.7) \quad \text{tr} P = \left(\frac{1}{2\pi}\right)^n \int \text{tr} p(\xi, x) d^n \xi d^n x$$

Note that P is a trace-class operator iff the order of its principal symbol is less or equal to $-1-\dim M$.

As an exercise, let us compute (3.6) when $M = S^1$ and P, Q are multiplication operators (infinitesimal gauge transformations). In that case the symbols are just smooth functions of the coordinate x on the circle. Now $\epsilon = \frac{\xi}{|\xi|}$ is a step function on the real line, its derivative is twice the Dirac delta function located at $\xi = 0$. It follows that the symbol of the commutator $\frac{1}{2}[\epsilon, P]$ is

$$(-i)\delta_\xi p'(x) + \frac{(-i)^2}{2!}\delta'_\xi p''(x) + \dots$$

Applying the formula (3.7) to (3.6) we get

$$\frac{1}{4}\text{tr} \epsilon[\epsilon, P][\epsilon, Q] = \frac{i}{2\pi} \int_{S^1} \text{tr} p'(x)q(x)dx,$$

where the trace under the integral sign is an ordinary matrix trace. If one feels uneasy with singular symbols, one can approximate ϵ by a differentiable function $\frac{\xi}{|\xi|+\lambda}$ and at the very end let $\lambda \mapsto 0$.

In the 3-dimensional case we have to insert $P = \theta(X; A), Q = \theta(Y; A)$ in (3.6). Using the asymptotic expansions for P and Q , $p = \sum p_{-k}(\xi, x)$ one has

$$(3.8) \quad c_3(X, Y; A) = \sum_{j,k} \text{tr} \left[\frac{\xi}{|\xi|}, p_{-j} \right] * q_{-k}$$

In fact, one needs to take into account only finite number of terms. The sum of terms with $j + k \geq 4$ is a coboundary of the 1-cochain

$$(3.9) \quad \sum_{k \geq 4} \text{tr} (\epsilon * \theta(X; A)_{-k})$$

Thus we may restrict the sum in (3.8) to indices $j + k < 4$. To take care of the infrared singularity in the integration in (3.7) we replace all denominators $|\xi|^{-k}$ by

$(|\xi| + \lambda)^{-k}$. One can then check by a direct computation that, modulo coboundaries, the result of the computation in (3.8) does not depend on the value of λ (i.e., one may take the limit $\lambda \mapsto 0$ in cohomology). The final result is in accordance with the cohomological [M, F-S], [M2], and perturbative arguments, [JJ],

$$(3.10) \quad c_3(X, Y; A) = \frac{1}{24\pi^2} \int_M \text{tr} A[dX, dY].$$

REFERENCES

- [A] H. Araki in: *Contemporary Mathematics*, vol. 62, American Mathematical Society, Providence (1987)
- [JJ] R. Jackiw and K. Johnson, Phys. Rev. **182**, 1459 (1969)
- [L] Lars-Erik Lundberg, Commun. Math. Phys. **50**, , 103 (1976)
- [M, F-S] J. Mickelsson, Commun. Math. Phys. **97**, 361 (1985); L. Faddeev, S. Shatasvili, Theoret. Math. Phys. **60**, 770 (1984)
- [M1] J. Mickelsson, Lett. Math. Phys. (1993)
- [M2] J. Mickelsson, *Current Algebras and Groups*, Plenum Press, New York and London (1989)
- [PS] A. Pressley and G. Segal, *Loop Groups*, Clarendon Press, Oxford (1986)